

## **Predictors of Failure in Reverse Obliquity Intertrochanteric Femur Fractures**

Christian A Gonzalez<sup>1</sup>, Noelle Lily Van Rysselberghe<sup>2</sup>, Mathew James Whittaker, Daniel Ngo, Harin Bhavin Parikh<sup>3</sup>, Juntian Wang, Abrianna Robles<sup>4</sup>, Andrea Helen Horne, Garrett Allen Cavanaugh, Garrett Esper, Arya Amirhekmat, Eleni Berhaneselase, Natalie Marie Marengi<sup>5</sup>, Marisa McDow, Christopher Gabriel Herbosa, Maricela Diaz, Uchechukwu Emili<sup>6</sup>, Zachary Lim, Dmitry Pokhvashchev, Aden Malik, Edmond F O'Donnell, Muhammad Umar Jawad<sup>7</sup>, Sean T Campbell, Milton Thomas M Little<sup>4</sup>, Walter W Virkus, Philipp Leucht, Matthew Robert Garner, Mark A Lee, John Alan Sclaro<sup>8</sup>, Marschall B Berkes, Saam Morshed, Stephen James Warner<sup>9</sup>, Paul William Perdue<sup>10</sup>, Eben A Carroll<sup>11</sup>, Justin Lucas<sup>12</sup>, Julius A Bishop, Lawrence Henry Goodnough, Michael J Gardner<sup>13</sup>

<sup>1</sup>University of Nevada, Reno School of Medicine, <sup>2</sup>Stanford University, <sup>3</sup>Cedars-Sinai Medical Center, <sup>4</sup>Cedars Sinai Medical Center, <sup>5</sup>Wake Forest, <sup>6</sup>OSU Orthopedics, <sup>7</sup>Samaritan Health, <sup>8</sup>University of California, Irvine - Orthopaedics, <sup>9</sup>McGovern Medical School At Uthealth, <sup>10</sup>VCU, <sup>11</sup>Wake Forest Univ School Of Medicine, <sup>12</sup>Santa Clara Valley Medical Center, <sup>13</sup>Stanford University Sugery

**INTRODUCTION:** Reverse obliquity (AO/OTA 31-A3) intertrochanteric (IT) femur fractures are associated with a high degree of patient morbidity and mortality. While negative effects of varus malreductions in proximal femur biomechanics are well known, the effect of coronal plane reduction on fixation failure in reverse obliquity IT fractures specifically has not been established. The purpose of this study was to assess the rates and risk factors for fixation failure, nonunion, and reoperation in reverse obliquity IT fracture treated with cephalomedullary nails (CMNs). We hypothesized the rate of fixation failure would be <5% when reduced with a postoperative neck shaft angle (NSA) >125°.

**METHODS:** Adults who sustained IT fractures (AO/OTA 31A1-A3) between 1/2014 and 5/2021 at thirteen level I trauma centers across the United States treated with CMNs with a minimum of 3 months follow up were included. Exclusion criteria included non-IT fractures and pathologic fractures. Initial differences in outcomes were analyzed with a chi-squared test. Backward elimination multivariable regression that accounted for age, sex, mechanism, fracture pattern, and acceptable reduction quality (defined as a postoperative NSA > 125°) was used to identify risk factors for adverse outcomes.

**RESULTS:** There were 2,132 intertrochanteric fractures identified during the study period, of which 305 (14.3%) were reverse obliquity fractures. Complications included 5 cases (1.6%) of fixation failure, 3 nonunions (1.0%), and 16 reoperations (5.2%). Age, sex, and mechanism of injury were not significantly associated with any outcomes. Compared to patients with adequate reduction, varus malreduction was significantly associated with fixation failure (5.5% vs. 0.8%,  $p = 0.014$ ), and reoperation (10.9% vs. 4.0%,  $p = 0.039$ ), but was not significantly associated with nonunion. Only varus malreduction (defined as a postoperative NSA  $\leq 125^\circ$ ) was a significant risk factor for fixation failure (OR 7.07 [95% CI 1.15-43.36],  $p = 0.035$ ) as well as reoperation (OR 2.90 [95% CI-1.01-8.36],  $p = 0.048$ ).

**DISCUSSION AND CONCLUSION:** Postoperative coronal neck shaft angle <125 degrees was associated with a greater than seven times increase in the odds of fixation failure and almost triple the odds of reoperation in reverse obliquity intertrochanteric femur fractures treated with cephalomedullary nails. Less than one percent of cases with adequate reduction had fixation failure, suggesting excellent results can be achieved in this morbid fracture pattern with proper reduction.