

# **Is it Healed? Clinical Measures of Fracture Healing Do Not Predict Unplanned Nonunion Surgery in Distal Femur Fractures**

Kelly Ann Lefavre<sup>1</sup>, Gerard Slobogean<sup>2</sup>, Peter Joseph O'Brien

<sup>1</sup>Vancouver General Hospital/ University of British, <sup>2</sup>University of Maryland School of Medicine

## **INTRODUCTION:**

Historically, measures of success after distal femur fracture fixation have relied on radiographic outcomes. Newer instruments bridge the gap between radiographic union and functional outcomes by assessing "clinical union." The Function IndeX for Trauma (FIX-IT) score quantifies clinical healing by aggregating assessments of weightbearing and fracture site pain. We sought to determine the relationship between clinical healing, radiographic healing, and traditional functional outcome measures.

**METHODS:** This secondary analysis of randomized controlled trial data included 167 adult patients with an OTA type 33A/C distal femur fracture recruited from 15 centers. Eligible fractures were treated with a bridging construct using a lateral locked plate and randomly assigned to either far cortical locking screws or standard screws. Outcomes were assessed at 6 weeks and 3, 6, and 12 months after surgery. Our primary comparison assessed time-specific correlations between clinical healing, measured by the FIX-IT score, and radiographic healing, measured by the radiographic union score for tibia (RUST). Secondarily, we assessed time-specific correlations between clinical healing and physical function, measured with the 36-item short-form health survey (SF-36). Finally, we constructed models to detect cut-points that optimized accuracy for predicting major reoperations using the RUST and FIX-IT scores at 6 weeks and 3 months.

## **RESULTS:**

Correlations between RUST and FIX-IT scores were negligible at 6 weeks ( $r=0.02$ ) and 3 months ( $r=0.12$ ), improved at 6 months ( $r=0.37$ ), but diminished at 12 months ( $r=0.19$ ). Correlations between FIX-IT and SF-36 Physical Component Scores were initially weak but improved to become moderately stronger ( $r=0.17$  at 6 weeks;  $r=0.39$  at 3 months;  $r=0.52$  at 6 months;  $r=0.59$  at 12 months). Cut-points for the RUST score of 6 at 6 weeks (AUC=0.74) and 9 at 3 months (AUC=0.90) showed reasonable accuracy in predicting major reoperation (0.75 and 0.80, respectively), with no improvement in performance with the addition of FIX-IT scores.

## **DISCUSSION AND CONCLUSION:**

Despite the importance of ensuring a distal femur fracture has clinically and radiographically healed, there is limited concordance between these assessments. Moreover, radiographic healing is more predictive of reoperations than the patient-centered focus on clinical fracture healing.