

# Timing of Radiographic Healing for Distal Femur Fractures Treated with an Intramedullary Nail

Dane Jensen Brodke, Nathan N O'Hara<sup>1</sup>, Sai Devana, Adolfo Hernandez, Cynthia Burke, Jayesh Gupta, Natasha S. McKibben<sup>2</sup>, Robert V O'Toole, John Morellato<sup>3</sup>, William Hunter Gillon, Murphy McGriff Walters<sup>4</sup>, Colby Barber<sup>5</sup>, Paul William Perdue<sup>6</sup>, Graham John Dekeyser, Lillia Steffenson<sup>7</sup>, Lucas Scott Marchand<sup>7</sup>, Marshall James Fairres<sup>8</sup>, Loren O Black, Zachary Mark Working, Erika Roddy, Ashraf N El Naga, Matthew Hogue<sup>9</sup>, Trevor Gulbrandsen<sup>10</sup>, Omar Hammad Atassi, Thomas W. Mitchell, Stephen J Shymon<sup>11</sup>, Christopher Lee<sup>12</sup>

<sup>1</sup>University of Maryland, <sup>2</sup>Oregon Health & Science University, <sup>3</sup>University of Mississippi Medical Center, <sup>4</sup>UMMC, <sup>5</sup>Colby Barber, <sup>6</sup>VCU, <sup>7</sup>University of Utah, <sup>8</sup>Harbor UCLA, <sup>9</sup>UIHC, <sup>10</sup>University of Iowa Hospitals, <sup>11</sup>Los Angeles County Harbor-UCLA Medical Center, <sup>12</sup>UCLA

## INTRODUCTION:

Modified Radiological Union Scale for Tibia (mRUST) scores have been found to be a reliable metric of radiographic healing in distal femur fractures treated with intramedullary nails. The purpose of this study was to descriptively profile mRUST scores in a large cohort of distal femur fractures and analyze predictors of timing of healing.

## METHODS:

A multicenter retrospective cohort study was undertaken. Patients presenting to one of 10 tertiary care referral centers with a distal femur fracture (OTA 33A or C) between 2012 and 2019, treated with an isolated intramedullary nail, were eligible for inclusion. Patients with less than 1 year of radiographic follow up were excluded unless they achieved radiographic union (defined as an mRUST of 12) or underwent unplanned reoperation to promote union prior to 1 year. mRUST scores collected at 3, 6, and 12 months were graphed for each patient (Figure). ROC curve analysis was utilized to identify the optimal cutoff in 3-month mRUST score that predicted reoperation to promote union. Multivariable models identified predictors of the timing of radiographic union and predictors of delayed progression.

## RESULTS:

A total of 157 fractures in 154 patients were included. Mean follow up was 17 months. In total, 118 fractures achieved radiographic union at a mean 6.2 months. The timing of radiographic union was predicted by female sex (1.2 months longer;  $p = 0.04$ ), tobacco use (1.3 months longer;  $p = 0.03$ ), open fracture (1.2 months longer;  $p = 0.04$ ), and the use of topical antibiotics (1.9 months longer;  $p = 0.03$ ). Delayed progression was predicted by chronic kidney disease (OR 13.7;  $p < 0.01$ ). A 3-month mRUST score of  $\leq 8$  predicted reoperation to promote union with a sensitivity of 88%, a specificity of 70%, a positive predictive value of 26%, and a negative predictive value of 98%.

## DISCUSSION AND CONCLUSION:

Distal femur fractures treated with intramedullary nail fixation heal radiographically at 6 months on average and have a 98% likelihood of treatment success if an mRUST threshold of  $\geq 9$  is surpassed at 3 months postoperatively. Factors associated with prolonged healing included female sex, tobacco use, open fracture, and the use of topical antibiotics. Surgeons can use this information to better understand the normal healing trajectory of distal femur fractures and stratify patients prognostically.

