

# **Do Patient or Surgical Factors Impact Outcomes after Traumatic Elbow Instability: Analysis of 201 Complex Elbow Fracture-Dislocations**

Christen E Chalmers, Amanda Tedesco, Jacquelyn Cruz<sup>1</sup>, Gerardo Sanchez-Navarro, Maike Van Niekerk<sup>2</sup>, Jacques Henri Hacquebord, Philip Lim<sup>3</sup>, Babar Shafiq<sup>4</sup>, Michael J Gardner<sup>5</sup>, Ranjan Gupta<sup>6</sup>

<sup>1</sup>Stanford Health Care, <sup>2</sup>Stanford University, <sup>3</sup>UC Irvine, <sup>4</sup>Johns Hopkins University, <sup>5</sup>Stanford University Sugery, <sup>6</sup>Univ of California Med Ctr

## **INTRODUCTION:**

Management of elbow instability remains challenging despite advancements in surgical technique and implant design. Restoration of stability to allow for early range of motion is paramount; yet which factors impact outcomes remains unclear. We hypothesized delayed intervention, combined ulnohumeral/radiocapitellar instability, and increased operative time would increase complications.

## **METHODS:**

We initiated an IRB-approved multicenter retrospective review of patients  $\geq 18$  years old who underwent surgical intervention for elbow fractures with concomitant instability between 2016-2022 from four level 1 trauma centers, excluding those with prior elbow surgery, pathologic fracture, intact joint stability, or  $< 3$  months follow-up. Patient inclusion was confirmed by fellowship-trained orthopedic surgeons after reviewing injury radiographs. Demographics, comorbidities, injury patterns, operative time, postoperative complications, and final ROM were recorded. Statistical analyses including rank-biserial correlations, Spearman's correlations, and Fisher's exact tests were performed using IBM SPSS software with significance level of  $p < 0.05$ .

## **RESULTS:**

A total of 201 patients [47.3% male; age mean (range)=51.3 (18-93)] met inclusion criteria with transolecranon fracture-dislocations as most common injury (32.7%). Most patients demonstrated both radiocapitellar and ulnohumeral instability (82.3%) and underwent surgical intervention at median of 5.5 days from injury. Post-operative complications included heterotopic ossification (HO) (34.3%), revision surgery (20.9%), neurological deficits (15.3%), surgical site infection (5.5%), and hardware failure (4.5%). Prolonged operative time was associated with increased incidence of revision surgery ( $r=0.301$ ,  $p=0.014$ ) and decreased elbow extension ( $r=0.469$ ,  $p<0.001$ ) and supination ( $r=-0.308$ ,  $p=0.033$ ). HO development – also associated with decreased elbow extension ( $r=0.327$ ,  $p=0.007$ ) and supination ( $r=-0.356$ ,  $p=0.013$ ) – was correlated with ligamentous repair ( $p=0.028$ ), but not isolated bony fixation procedures or revision surgery. Time from injury to surgery, injury pattern, and radiographic instability were not significantly correlated with complications.

## **DISCUSSION AND CONCLUSION:**

Prolonged operative time was associated with reduced elbow mobility and greater revision surgery incidence. Ligamentous repair to restore joint stability paradoxically increased the risk of HO and stiffness. The data supports that surgical decision-making, not patient's injury pattern, markedly impacts outcomes. This data provides impetus for greater efforts to develop standardized algorithms for the surgical management of complex traumatic elbow instability.