



AMERICAN ACADEMY OF  
ORTHOPAEDIC SURGEONS

# 2023 Annual Meeting

March 7 – 11, Las Vegas, Nevada

**Session: V**

**Session Title:** Disaster-Plasty: A Case-Based Guide to Leveraging Trauma and Arthroplasty Skills for the Extreme Challenges

**Session Type:** Symposium

**Location:** Room 4303

**Date & Time:** 03-10-2023, 11:00 am - 12:30 pm

**INSTRUCTORS WHO CONTRIBUTED TO THIS HANDOUT:** as of 1/24/2023

**Moderator(s):**

George J. Haidukewych, MD, FAAOS

**Faculty:**

Derek J. Donegan, MD, FAAOS

Elizabeth Gausden, MD

Erik Kubiak, MD, FAAOS

Gwo-Chin Lee, MD, FAAOS

Brett R. Levine, MD, FAAOS

Frank A. Liporace, MD, FAAOS

Theodore T. Manson, MD, FAAOS

Chloe Scott, BSc

Richard S. Yoon, MD

Brandon J. Yuan, MD, FAAOS

The Academy reserves any and all of its rights to materials presented at the Annual Meeting. Reproductions of any kind, by any person or entity, without prior written permission from the Academy are strictly prohibited.

**AAOS 2023--Disaster-plasty: A case-based guide to leveraging trauma and arthroplasty skills for the extreme challenges**

**Moderator: George Haidukewych, M.D. (Orlando Health)**

**OUTLINE (90 minutes)**

Introduction- GH (5 minutes)

1. Fractured... and Infected: Dealing with infections following periprosthetic fractures (Chloe Scott, M.D.) (7 minutes)
2. Periprosthetic femur fracture nonunion: What do I do now? (Brandon Yuan, M.D.) (7 minutes)
3. Periprosthetic Fractures of the Femur with Massive Bone Loss: Managing with Revision Arthroplasty (Brian Gladnick, M.D.) (7 minutes)
4. Periprosthetic acetabular fractures: Approach and timing (Richard Yoon, M.D.) (7 minutes)
5. Periprosthetic fractures of the tibia: use of the extreme tibial nail and when to plate (Frank Liporace, M.D.) (7 minutes)
6. Malunion and arthroplasty: Addressing alignment through revision
7. (Elizabeth B. Gausden, M.D.) (7 minutes)
8. Revision TKA in setting of massive osteolysis: Cones have changed the game (Gwo-Chin Lee, M.D.) (7 minutes)
9. Bridge plating to prevent inter-prosthetic femur fractures: Call me now or call me later! (Eric Kubiak, M.D.) (7 minutes)
10. Use of distal femoral replacement for periprosthetic distal femur fractures: My tips to make this last (Brett Levine, M.D.) (7 minutes)
11. Extreme nailing of periprosthetic distal femur fractures (Derek Donegan, M.D.)
12. Rapid Fire Case-Based Examples (George Haidukewych, M.D. & Charlie Hannon, M.D.) (10 minutes)

## **Fractured... and Infected: Dealing with infections following periprosthetic fractures**

Chloe Scott, MD, MSc

Edinburgh Orthopaedics, Royal Infirmary of Edinburgh, UK

### Definition:

- Rate of fracture-related infection (FRI) very variable: 3 – 20%
- Definitely infected: wound breakdown and purulent drainage
- Possibly infected: Erythema, warmth and swelling. Also present in aseptic non-union.

### Diagnosis:

- *Blood tests:* CRP and blood cultures.
- *Imaging:* Fracture union, implant loosening and osteomyelitis (periosteal reaction) may be evident on plain radiographs but may require CT which will also identify sequestrations and associated collections. Collections/effusions should be aspirated +/- ultrasound guidance.
- *Microbiology:* aspirates and deep tissue samples essential for microbiology and histology. Do not send wound edges/sinus tracts due to contamination with multiple organisms.

### Avoiding infection

#### Soft tissue management:

- *Timing of surgery:* the majority of PPFs are low energy and benefit from prompt fixation. Some high energy injuries with significant soft tissues injury (eg. button holing through knee capsule) may require surgical delay.
- *Approach:* haemostasis with sequential identification and tying off of perforating vessels minimizes blood loss, haematoma formation and postoperative anaemia
- *Skin closure:* consider deep tension nylon mattress suture closure in fragile skin/high BMI
- *Negative pressure dressings:* where there is concern about the potential for wound leakage.
- *Resting:* splinting knees in extension for 1-2 weeks until the skin has healed or using negative pressure dressings can be useful where skin is fragile or multiply operated

### Antibiotics:

- Extended antibiotic prophylaxis (for 3-7 days), intra-incisional Vancomycin; antibiotic cement.

### Medical management:

- *Anticoagulation:* DOACs can cause wound leakage in elderly. Use LMWH in preference until skin has healed.
- *Optimize medical management* – manage diabetes and anaemia. Orthogeriatric and microbiology/infectious diseases multidisciplinary care.

### Management of PPF infection:

- Antibiotic suppression until fracture union AND surgical eradication of the biofilm
- Fracture stability is essential for infection eradication.
- Mean time to periprosthetic femoral fracture union is ~6 months - osteomyelitis can occur while waiting for union.
- Staph aureus and coagulase negative staph are most common organisms in FRI.
- Treatment strategy (Eradication Vs long term suppression) determined by fracture pattern, organism, and host.
- Eradication requires a stable construct, a good soft tissue envelope and an effective surgical debridement.
- Scenario 1: Bone preservation and infection eradication
  - Requires good host and easy to treat infection (sensitive single organism)
  - Debride and sample, suppress, obtain union, revise (in 1 or 2 stages).

- Scenario 2: Non-bone preserving infection eradication
  - Requires good host and easy to treat organism
  - Radical debridement, single stage endoprosthetic replacement
- Scenario 3: Long term suppression
  - When the implant must be retained (fracture not united) OR difficult to treat organism (multiresistant/polymicrobial) OR unfit host.
  - Washout, debridement, and sampling followed by antibiotic suppression +/- sinus formation. As implant is retained biofilm cannot be eradicated.
  - May be needed in frail elderly patients regardless of organism if too frail for extensive revision surgery. This approach may also be required in less frail patients with resistant organisms that are difficult to treat or polymicrobial infection.

**Periprosthetic femur fracture nonunion: What do I do now? (Brandon Yuan, M.D.) (7 minutes)**

- Walk step-wise through the nonunion work up process with case examples:
  - Step 1: Why didn't it heal?
    - Patient issue?
    - Infection? (I will just mention given Dr. Scott going over this just before)
    - Biology?
    - Technical/Mechanical Issue?
  - Step 2: Can that problem be corrected, and most importantly, can the patient tolerate it?
    - How to correct technical/mechanical issues
    - How to add biology
    - Don't forget to assess if arthroplasty is worth saving
  - Step 3: Decide – Revision ORIF vs. Revision arthroplasty
    - What are the advantages and disadvantages of each approach?
  - Step 4: Execute
    - Correcting the technical/mechanical issue - Case example of revision ORIF
    - Bypassing the biologic or patient issue – Case example of revision arthroplasty
    - When do you need to do both? – Case example of ORIF + revision (examples of femoral nonunions above unstable CR TKAs, I have some video of this)

## **Periprosthetic Fractures of the Femur with Massive Bone Loss: Managing with Revision Arthroplasty**

**(Brian Gladnick, M.D.) (7 minutes)**

### **Background:**

Periprosthetic fractures of the femur are a devastating complication after total hip arthroplasty and are associated with significant morbidity. Risk of complications such as fracture, instability, and infection are increased in these high-risk patients, and mortality after revision may be as high as 21% at one year post-operatively [1].

Perhaps one of the most challenging aspects of treating these injuries is dealing with the inherent bone loss that is typically seen in the revision scenario, but can be compounded in the periprosthetic fracture setting [2]. Patients are typically older, with poor bone quality and low bone mineral density. Furthermore, comminution at the fracture site or bone loss that occurs during cement removal may compound these reconstructive challenges.

One of the most important components of pre-operative planning for revision THA of a periprosthetic femur fracture is assessing where the best residual bone exists, and templating a reconstruction that maximizes the chance of getting durable mechanical fixation with a stable fracture construct [3]. The Vancouver classification is the most well-known and useful classification in this scenario and provides an algorithm for fixation that is matched to the patient's fracture type. Vancouver B2 fractures are fractures that occur around the femoral component, with coincident debonding of the implant from bone, cement, or both. In Vancouver B3 fractures, the stem is similarly loose, but with the added complication of poor proximal bone stock. While Vancouver B2 fractures may commonly be reduced anatomically and revised with either a longer primary or revision stem, Vancouver B3 fractures often are more challenging to reduce, and may require even longer modular stems with or without bone grafting in order to achieve a stable construct, ideally with at least 4cm of stem-cortical contact length [4,5].

Finally, intraprosthetic fractures below the tip of a well-fixed stem can be a unique challenge due to the difficulty of achieving either proximal or distal fixation, and may require longer revision constructs to bridge the fracture zone, particularly when prior plating has been unsuccessful [6].

### **Case 1:**

Vancouver B3 fractures may occur in situations in which a previously loose stem has caused progressive proximal bone loss, ultimately resulting in a fracture due to the weakened bone, mechanical fall, or a combination of both. Osteolysis from polyethylene wear may influence the loss of proximal bone stock, and pre-operative radiographs should be scrutinized for eccentric wear of the femoral head.

In the present case, a loose cemented stem in a 79-year-old female has caused progressive proximal bone loss, with a capacious proximal canal and cortical thinning, especially posteriorly. Pre-operative plan was to bypass this area of poor proximal bone with a modular, splined tapered stem, achieving at least 4cm of cortical contact distally. Intra-operative clinical photographs show that the stem was grossly loose, and was removed through a direct anterior approach with its cement mantle intact. Reduction was achieved with luque wires and a turkey-claw fracture clamp, which allows for stable fracture reduction while reaming. A wire has been placed below the fracture site to prevent distal propagation. Post-op films demonstrate reduction of the fracture, which has been bypassed by a modular splined tapered stem with at least 4cm of cortical contact distally. Note the proximal bone loss and thinned cortices on the post-operative view.

### **Case 2:**

In the setting of conversion arthroplasty, the first step in pre-operative planning should always be to

rule out infection. This is particularly concerning in the setting of femoral nonunion, where the rate of indolent infection may be as high as 21-44%, and the offending organism is most often *Staphylococcus* species [7]. Patients should be questioned specifically about infection history surrounding the previous fracture surgery, including wound healing issues, prolonged drainage, extended use of antibiotics, or return trips to the OR for irrigation and debridement. Laboratory workup includes ESR and CRP; if these labs are elevated or if there is any clinical suspicion for infection, aspiration and culture should be performed. If radiology aspiration is not feasible, a staged procedure including removal of hardware may be performed in order to obtain samples of the fracture site and hardware for culture to rule out infection prior to definitive arthroplasty. Once infection has been definitely ruled out, arthroplasty may proceed; however care must be taken to address any bone defects, screws holes, or other stress risers which may exist after removal of the femoral hardware. If the femoral architecture is relatively preserved, reconstruction may be accomplished with a primary-type stem, but in cases of nonunion, malunion, or severe proximal bone defect, it may be necessary to bypass the deficient area with a distally engaging revision stem.

In the present case, a 68-year-old female presents with an extensive surgical history involving her left femur, including multiple revision ORIF procedures for nonunion and ultimately requiring a vascularized fibular autograft to achieve femoral union. Her pre-operative ESR and CRP were elevated. Because of the likely low utility of IR aspiration of the fracture site, she instead underwent a staged procedure including removal of multiple proximal screws in order to culture both the hardware and screw tracts prior to definitive arthroplasty. While cultures were all negative for infection, unfortunately she suffered a fall from her wheelchair four weeks out from hardware removal, resulting in a transverse fracture through one of the residual screw tracts, greatly increasing the complexity of the reconstruction.

A two-incision approach is planned, as seen in the preoperative clinical photograph. The fracture is provisionally reduced with a plate applied via a lateral approach to the lateral aspect of the femur via cables. Definitive stability is achieved by placing a modular splined-tapered stem through a direct anterior approach. The tapered, splined stem design provides axial and torsional stability, essentially acting like an intramedullary nail. Proximally, a modular cone body is mated to the distal stem component, completing the femoral component of the arthroplasty.

### Case 3:

Vancouver C periprosthetic fractures are located distal to the tip of a well fixed femoral stem, and are usually amenable to open reduction and plating [4]. However, the presence of the femoral stem in the canal of the femur increases the complexity of the biomechanical environment of the fracture. A stress riser at the tip of the stem, particularly in a stiff, distally fitting stem, may significantly impair fracture healing. Furthermore, proximal fixation is compromised due to the presence of the femoral stem. In the setting of a concomitant knee replacement, distal fixation may also be compromised; these interprosthetic femur fractures are particularly challenging and have a revision rate as high as 24% [6].

The present case is a 78-year-old femur with a previous left THA and left TKA, who presented with a short oblique Vancouver C interprosthetic fracture below the tip of an SROM stem. This was initially treated with open reduction and lateral plating. Due to limited screw options proximally with the canal completely filled by the distally-reamed SROM stem, proximal fixation was limited to using a combination of cables and unicortical locking screws. The patient did well clinically and was fully weight bearing by three months post-operatively. However, at one year post-operative, she re-presented acutely with a hypertrophic varus malunion at the fracture site and fracture of the lateral plate.

In order to stabilize her fracture site and promote bony healing, the SROM stem was revised to a longer diaphyseal-engaging stem that bypasses the fracture site, acting as an intramedullary nail. A lateral plate was then re-applied, spanning the entire femur, and a fibular strut graft was cabled

anteriorly across the fracture site. The patient went on to achieve full radiographic union with this construct.

#### Case 4:

Interprosthetic fractures in which the patient has undergone previous revision arthroplasty are a unique reconstructive challenge. The presence of a revision implant (including stems, cones, sleeves, or other internal fixation) often makes revision surgery with typical revision implants difficult and sometimes impossible, because the patient may lack enough residual bone for fixation either proximally or distally. In these patients, bypassing the entire construct with a total femur replacement may be the only durable reconstructive option. Patients should be adequately counseled pre-operatively, as these operations have a high complication rate, and post-operative functional status may be limited by the extensive dissection and bone resection often required in such procedures [8].

In the present case, a 61 year-old-male presents with a right femur interprosthetic fracture between a total hip replacement and a distal femoral replacement, which had already failed two attempts at ORIF. The presence of a distal femoral replacement prevents the possibility of spanning the fracture with a long-stemmed THA construct, hence the decision was made to move forward by spanning the entire femur with a total femur replacement. Due to the patient's young age and prior functional status, the implant of choice was a bone-conserving intramedullary total femur. This device acts as an intramedullary nail spanning the fracture zone, but is a distal femur endoprosthesis at the knee, and proximally mates with a THA cone body, similar to reconstruction with any modular stem. A trochanteric claw plate which locks into the cone body was used to stabilize a fracture of the greater trochanter. Six year follow up films demonstrate durability of the endoprosthetic construct and healing of the diaphyseal nonunion.

#### References

1. Maradit Kremers H, Abdel MP, Ransom JE, Larson DR, Lewallen DG, Berry DJ. Mortality After Periprosthetic Femur Fractures During and After Primary and Revision Total Hip Arthroplasty. *J Am Acad Orthop Surg*. 2019 May 15;27(10):375-380. doi: 10.5435/JAAOS-D-17-00902.
2. Menken LG, Rodriguez JA. Femoral revision for periprosthetic fracture in total hip arthroplasty. *J Clin Orthop Trauma*. 2020 Jan-Feb;11(1):16-21. doi: 10.1016/j.jcot.2019.12.003. Epub 2019 Dec 14.
3. Rodriguez J.A. Preoperative planning and surgical technique in the management of periprosthetic femoral fractures using a tapered modular fluted prosthesis with distal fixation. *Oper Tech Orthop*. 2009;19:137-142
4. Naqvi G.A., Baig S.A., Awan , N. Interobserver and intraobserver reliability and validity of the vancouver classification system of periprosthetic femoral fractures after hip arthroplasty. *J Arthroplast*. 2012;27:1047-1050.
5. Ioannis Gkiatas 1, Michael-Alexander Malahias 1, William Xiang 1, Kathleen N Meyers 2, Lisa A Torres 2, T David Tarity 1, Jose A Rodriguez 1, Mathias P Bostrom 1, Timothy M Wright 2, Peter K Sculco 1. How Does Contact Length Impact Titanium Tapered Splined Stem Stability: A Biomechanical Matched Pair Cadaveric Study. *J Arthroplasty*. 2021 Sep;36(9):3333-3339. doi: 10.1016/j.arth.2021.04.013. Epub 2021 Apr 22
6. Tibbo ME, Limberg AK, Gausden EB, Huang P, Perry KI, Yuan BJ, Berry DJ, Abdel MP. Outcomes of operatively treated interprosthetic femoral fractures. *Bone Joint J*. 2021 Jul;103-B(7 Supple B):122-128.
7. Hackl S, Trenkwalder K, Militz M, Augat P, Stuby FM, von Rden C. Infected nonunion: diagnostic and therapeutic work-up. *Unfallchirurgie (Heidelb)*. 2022 Aug;125(8):602-610. doi: 10.1007/s00113-022-01204-w. Epub 2022 Jun 24.
8. Rao BM, Stokey P, Tanios M, Liu J, Ebraheim NA. A systematic review of the surgical outcomes of interprosthetic femur fractures. *J Orthop*. 2022 Jul 31;33:105-111. doi: 10.1016/j.jor.2022.07.013. eCollection 2022 Sep-Oct.



**Periprosthetic acetabular fractures: Approach and timing**  
**(Richard Yoon, M.D.) (7 minutes)**

- I.) Epidemiology
- II.) Risk Factors
- III.) Classification
- IV.) Treatment Options & When to Enlist Them
  - a. Non-operative management
  - b. Revision style jumbo cup as a plate
  - c. Plate plus multi-hole cup
  - d. Cup-cage
  - e. Triflange

**Malunion and Arthroplasty: Addressing alignment through revision**  
**Elizabeth B. Gausden, M.D., M.P.H.**

- I. Causes of extra-articular deformity around TKA
  - a. Congenital abnormalities
  - b. Posttraumatic malunion
  - c. Previous osteotomies
- II. Evaluation
  - a. Long leg hip to ankle radiographs
  - b. Dedicated knee films likely to miss extra-articular deformity
  - c. CT scan useful to assess for union in posttraumatic cases
  - d. Previous operative reports for assistance with hardware removal
- III. Planning
  - a. Assess leg length discrepancy
  - b. Intraarticular correction
    - i. If extra-articular deformity is <20 degrees in the coronal plane on femoral side or <30 degrees on the tibial side or when the CORA is located outside the metaphyseal region
  - c. More severe deformities, consider extra-articular osteotomy
    - i. If the intraarticular correction will compromise ligamentous attachments
    - ii. Staged technique
- IV. Cases
  - a. Malunion of periprosthetic distal femur fracture
  - b. Nonunion of a proximal tibial fracture with severe valgus deformity

**References**

Sculco PK, Kahlenberg CA, Fragomen AT, Rozbruch SR. Management of Extra-articular Deformity in the Setting of Total Knee Arthroplasty. J Am Acad Orthop Surg. 2019 Sep 15;27(18):e819-e830. doi: 10.5435/JAAOS-D-18-00361. PMID: 30624304.

Wang JW, Wang CJ. Total knee arthroplasty for arthritis of the knee with extra-articular deformity. J Bone Joint Surg Am. 2002 Oct;84(10):1769-74. doi: 10.2106/00004623-200210000-00005. PMID: 12377906.

## **Revision TKA in the Setting of Massive Osteolysis: Metaphyseal Cones and Sleeves have Changed the Game- Gwo-Chin Lee, MD**

Introduction: Improvements in surgical technique, prosthesis design and materials have not eliminated failures the need for revision surgery. Severe bone loss is a principal obstacle to implant longevity in revision arthroplasty and therefore a systematic approach to management of bone loss during revision surgery is critical to maximizing function and minimizing failures. In recent years, our understanding of the importance of maximizing fixation in the metaphysis to the stability of the revision TKA construct has greatly improved. Cones and sleeves have changed the way we approach case with severe bone loss and improved long term results. Below are the reasons on how cones/sleeves have changed the game.

- Cones and sleeves help restore joint line.
  - Restoration of the joint line improves knee joint stability and knee joint kinematics
    - Minimizes joint line elevation on the femoral side that can lead to mid-flexion instability [1].
    - Minimizes the need to downsize the tibial component and the need to offset a corresponding smaller femoral component in order to restore posterior condylar offset which is critical to flexion gap stability [2].
- Cones and sleeves improve axial and rotational stability of the revision TKA implant.
  - Reduces the need for longer, diaphyseal engaging stems and allows the use of shorter cemented stems [3].
    - Advantages of shorter cemented stems
      - Antibiotic delivery throughout the entire revision construct
      - Reduces the risk of anatomic conflict and the need for offset stems
      - Lowers the risk of end of stem pain
- Cones and sleeves improve fixation in patients requiring hinged knee or distal femoral replacement prosthesis.
  - Improves rotational stability of the construct and reduces the risk of component loosening [4].

### **REFERENCES**

1. Evangelista PJ, Laster SK, Lenz NM, Sheth NP, Schwarzkopf R. A Computer Model of Mid-Flexion Instability in a Balanced Total Knee Arthroplasty. J Arthroplasty. 2018 Jul;33(7S):S265-S269.
2. Kannan A, O'Connell RS, Kalore N, Curtin BM, Hull JR, Jiranek WA. Revision TKA for Flexion Instability Improves Patient Reported Outcomes. J Arthroplasty. 2015 May;30(5):818-21.
3. Alipit V, Kirk A, Scholl D, Schmidig G, Springer BD, Lee GC. Micromotion Analysis of Various Tibial Constructs in Moderate Tibial Defects in Revision Total Knee Arthroplasty. J Arthroplasty. 2021 Jan;36(1):362-367.e1.
4. Cottino U, Abdel MP, Perry KI, Mara KC, Lewallen DG, Hanssen AD. Long-Term Results After Total Knee Arthroplasty with Contemporary Rotating-Hinge Prostheses. J Bone Joint Surg Am. 2017 Feb 15;99(4):324-330.

*Use of distal femoral replacement (DFR) for periprosthetic distal femur fractures: My tips to make this last.*

Brett Levine, MD, MS  
Professor of Orthopaedics  
Rush University Medical Center

Epidemiology:

- Defined as fractures of the femur occurring within 15cm of the joint line or 5cm from an IM stem
- Periprosthetic fracture rates after primary TKA range from 0.3% to 5.5%
- Risk factors—osteoporosis, stress mismatch, component malalignment, notching, length of time the prosthetic is in situ

Classification:

- Multiple classification systems exist (Lewis and Rorabeck; Su et al; Chen et al; DiGioia and Rubash; Neer et al)
- Bottom line—Displaced v. nondisplaced; implant stability (loose v. stable), location of fracture

Indications:

- Distal segment does not offer enough support for fixation
  - Too small a segment
  - Poor bone quality in remaining segment (osteolysis, stress shielding)
- Concomitant injuries that require immediate weightbearing
- Poor function of prior TKA

Technique:

- Can use old midline incision depending on the extent of the fracture with a medial parapatellar approach
- Swashbuckler incision and lateral approach
- Approach through the fracture site, stay subperiosteal and remove the distal femur—release all attachments
  - Usually start medially or laterally and elevate collateral, then work thru notch area and fracture to the opposite side of the femur

Tips:

- Measure your length roughly before removing the distal end of the femur
- Make a perpendicular cut to the anatomic axis of the femur (if they have a planar flatten this surface)
- If the distal femur is patulous then will press-fit cone and cement thru cone—use cement restrictor
- Assess hip and make sure there is not a prosthesis there already or that there is no DJD present in the hip
- If the hip is arthritic or there is limited femoral fixation that can be obtained can move to a staged or single surgery IM total femur
- In younger patients will use a press-fit, on-growth stem
- Make sure limb lengths are close by checking with other leg
- Assure femur is brought down so joint line is restored based on prior radiographs and the patella height on the femur
- Make sure adequate rotation is maintained on the femur—can use linea aspera, but always cheat a little more rotation than you think you need.
- Make sure you have secure tibial fixation (all three zones)
- Check and recheck patella tracking
- Close down the dead space and use a drain during closure
- +/- NPWD, ace wrap for compression and often immobilize for 10-14 days for wound and capsular healing

Extreme nailing of periprosthetic distal femur fractures (Derek Donegan, M.D.)

1. Treatment options
  - a. Setting yourself up for success

2. Why to nail?
  - a. Biomechanical advantages
  - b. Key concepts for successful nailing
3. Antegrade IMN
  - a. When
  - b. Why
  - c. How
4. Retrograde IMN
  - a. When
  - b. Why
  - c. How
5. Tips and Tricks
  - a. Density of fixation distally
  - b. TKA Poly management
6. Conclusion